arXiv:cond-mat/9801272v2 [cond-mat.mes-hall] 12 Nov 1998

Quasi-particles in fractional quantum Hall effect edge theories

R. A. J. van Elburg and K. Schoutens
Van der Waals-Zeeman Institute and Institute for Theoretical Physics
University of Amsterdam, Valckenierstraat 65, 1018 XE Amsterdam, The Netherlands
(February 1, 2008)

We propose a quasi-particle formulation of effective edge theories for the fractional quantum Hall effect. For the edge of a Laughlin state with filling fraction $\nu = \frac{1}{m}$, our fundamental quasi-particles are edge electrons of charge $-e$ and edge quasi-holes of charge $+\frac{e}{m}$. These quasi-particles satisfy exclusion statistics in the sense of Haldane. We exploit algebraic properties of edge electrons to derive a kinetic equation for charge transport between a $\nu = \frac{1}{m}$ fractional quantum Hall edge and a normal metal. We also analyze alternative 'Boltzmann' equations that are directly based on the exclusion statistics properties of edge quasi-particles. Generalizations to more general filling fractions (Jain series) are briefly discussed.

05.30.-d, 05.70.Ce, 11.25.Hf, 73.40Hm

Contents

I Introduction and summary **2**

II Exclusion statistics **3**
A Exclusion statistics and the fqHe . . . 3
B Exclusion statistics in CFT . . . 4

III Quasi-particles for the $\nu = \frac{1}{m}$ fqHe edge **4**
A Quasi-hole states . . . 5
B Edge electron states . . . 6
C Duality . . . 7
D The full spectrum . . . 8

IV Equilibrium quantities **9**
A Specific heat . . . 9
B Hall conductance . . . 10

V Jack polynomials and beyond **11**
A The operator H_{CS} . . . 11
B Norms and form factors . . . 12

VI Transport properties **13**
A Kinetic equation for inter edge transport . . . 14
B Interpretation in terms of exclusion statistics . . . 15

VII Conclusions **17**

APPENDIXES **18**

A Composite edges: Jain series **18**

B Form factor for general m **19**

I. INTRODUCTION AND SUMMARY

Low energy excitations over (fractional) quantum Hall effect (qHe) ground states are localized near the edge of a sample. Certain aspects of qHe phenomenology can therefore be captured by an *effective edge theory.* The unusual properties (notably, fractional charge and statistics) of bulk excitations over a fractional qHe (fqHe) ground state carry over to analogous properties of the fundamental edge excitations.

In the existing (theoretical) literature on qHe edge phenomena, the fundamental edge quasi-particles have not played an important role. Most, if not all, results have been obtained by exploiting bosonization schemes. For the analysis of edge-to-edge tunneling experiments, a combination of bosonization with techniques from integrable field theories has led to exact results for universal conductance curves[1,2]. What has been missing until now is a description of edge-to-edge transport phenomena directly in terms of excitations that are intrinsic to an edge in isolation. The authors of[3] have explored such a picture and have derived a number of strong-coupling selection rules for the scattering of edge excitations.

In this paper we take up the challenge of reformulating effective qHe edge theories directly in terms of a set of fundamental edge excitations. For the principal Laughlin states at filling fraction $\nu = 1/m$, we select the *edge electron* (of charge $-e$) and the *edge quasi-hole* (of charge $+e/m$) as the fundamental excitations. (The reason for this asymmetric choice will become clear.) An important complication is then that these fundamental quasi-particles are *not* fermions but instead obey fractional statistics (in a sense to be explained below). A large part of this paper will be devoted to working out the consequences of these unusual statistics.

Following Wen[4], we shall assume that the edge theory for a $\nu = 1/m$ fractional qHe state (Laughlin state) takes the form of a chiral Luttinger Liquid (see[5] for a further justification of this description). The bosonic description of such a theory is centered around the neutral charge density operators $n_q = \frac{1}{\sqrt{m}}(\partial\varphi)_q$, which satisfy a $U(1)$ affine Kac-Moody algebra

$$[n_q, n_{q'}] = \frac{1}{m}\, q\, \delta_{q+q'} \ . \tag{I.1}$$

The vertex operators

$$\Psi_{\rm e}(z) = \, : \exp(-i\sqrt{m}\varphi) : (z) \ , \qquad \Psi_{\rm qh}(z) = \, : \exp(i\frac{1}{\sqrt{m}}\varphi) : (z) \tag{I.2}$$

have charge $-e$ and $+\frac{e}{m}$, respectively, and have been identified as the edge electron and edge quasi-hole. They are the direct analogues of the bulk (Laughlin) quasi-particles, an important distinction being that the edge excitations are gap-less and have linear dispersion.

The first indication for the non-trivial statistics of the operators (I.2) comes from the Operator Product Expansions (OPE)

$$\Psi_{\rm e}(z)\Psi_{\rm e}(w) = (z-w)^m \, [\Psi'(w) + \ldots] \ , \qquad \Psi_{\rm qh}(z)\Psi_{\rm qh}(w) = (z-w)^{\frac{1}{m}} \, [\Psi''(w) + \ldots] \tag{I.3}$$

where $\Psi'(w)$ and $\Psi''(w)$ are operators of charge $-2e$ and $+\frac{2e}{m}$, respectively. For m an odd integer, the right hand side of the first OPE picks up a minus sign under the exchange $z \leftrightarrow w$, in correspondence with the required antisymmetry of the Laughlin wave functions. The second relation features a fractional power of $(z-w)$, which shows that the quasi-hole operator has fractional 'exchange statistics'.

For $\nu = 1$ the operators $\Psi_{\rm e}$ and $\Psi_{\rm qh}$ are fermionic and the edge theory is simply a theory of free fermions. The exchange statistics of $\Psi_{\rm e}$ and $\Psi_{\rm qh}$ at $\nu = 1/m$ clearly signal a deviation from free fermion behavior, but they are hardly helpful for the purpose of setting up a quasi-particle formalism that mimics the free fermion treatment of the $\nu = 1$ edge. We shall here argue that a much more convenient point of view is that of the 'exclusion statistics' properties of $\Psi_{\rm e}$ and $\Psi_{\rm qh}$.

In recent paper[6], one of us proposed a method to associate exclusion statistics to quasi-particles for Conformal Field Theory (CFT) spectra. We shall here show that, when applied to the $\nu = 1/m$ edge theory excitations, this method gives ideal fractional exclusion statistics (in the sense of Haldane[7]) with $g = m$ for edge electrons and $g = \frac{1}{m}$ for edge quasi-holes. We also find that the edge electrons and edge quasi-holes can be viewed as independent excitations, in the sense that there is no mutual exclusion between the two.

Our program in this paper is then (1) to establish the exclusion statistics properties of $\Psi_{\rm e}$ and $\Psi_{\rm qh}$, and (2) to apply them to both equilibrium and transport properties of these edges. As for transport, we shall focus on the set-up of the experiment by Chang et al.[8], where electrons are allowed to tunnel from a normal metal into the edge of a $\nu = 1/3$ fqHe edge. We shall use algebraic properties of $\nu = 1/3$ edge electrons to write an exact kinetic equation for the perturbative I-V characteristics for this system, reproducing the result obtained by other methods. Interestingly, the relevant algebraic properties derive from the so-called $N = 2$ superconformal algebra, which has been well-studied in the context of String Theory. We shall also study 'naive' Boltzmann equations that are based on the exclusion statistics properties of the edge quasi-particles. While the latter equations are not exact, we shall argue that they can be used as the starting point in a systematic approximation to the exact transport results. These results then are of general importance, as they illustrate the possibilities and limitations of the concept of exclusion statistics in the analysis of non-equilibrium physics.

The observations made here are easily generalized to composite edges, related to hierarchical fqHe states, in particular those of the Jain series with $\nu = \frac{n}{np+1}$. For the Jain series edge theories, two natural pictures emerge. In the first picture, the edge quasi-particles satisfy Haldane's exclusion statistics with G-matrix equal to K^{-1}, where K is the topological order matrix of the bulk fqHe state. In the second picture one decouples one charged mode from $n-1$ neutral modes. A possible quasi-particle basis then consists of a single charged Haldane g-on and a collection of n neutral quasi-particles that are related to parafermions in the sense of Gentile.

This paper is organized in the following way. In section 2 we discuss exclusion statistics and indicate the applications to fqHe states and to CFT spectra. In section 3 we discuss in some detail how a quasi-particle basis for $\nu = \frac{1}{m}$ fqHe edge states is obtained and how that leads to an assignment of exclusion statistics parameters. In section 4 we explain how equilibrium properties are obtained in a quasi-particle approach. In section 5 we further study the quasi-particle bases and make the link with Calogero-Sutherland quantum mechanics and Jack polynomials. In section 6 we study charge transport between a normal metal and a $\nu = \frac{1}{m}$ fqHe edge in terms of kinetic equations that are based on our quasi-particle formalism. The appendix A describes the extension of our quasi-particle formalism to filling fractions in the Jain series, while appendix B contains explicit results for an important quasi-particle form factor.

II. EXCLUSION STATISTICS

In his by now famous 1991 paper[7], Haldane proposed the notion 'fractional exclusion statistics', as a tool for the analysis of strongly correlated many-body systems. The central assumption that is made concerns the way a many-body spectrum is built by filling available one-particle states. In words, it is assumed that the act of filling a one-particle state effectively reduces the dimension of the space of remaining one-particle states by an amount g. The choices $g = 1$, $g = 0$ correspond to fermions and bosons, respectively. The thermodynamics for general 'g-ons', and in particular the appropriate generalization of the Fermi-Dirac distribution function, have been obtained in[9,10,11,12]. The so-called Wu equations[9]

$$n_g(\epsilon) = \frac{1}{[w(\epsilon) + g]} \qquad \text{with} \qquad [w(\epsilon)]^g [1 + w(\epsilon)]^{1-g} = e^{\beta(\epsilon-\mu)} \tag{II.1}$$

provide an implicit expression for the 1-particle distribution function $n_g(\epsilon)$ for g-ons at temperature T and chemical potential μ. It has been demonstrated that fractional exclusion statistics are realized in various models for Quantum Mechanics with inverse square exchange[7,13,14] and in the anyon model in a strong magnetic field[15,9].

A. Exclusion statistics and the fqHe

A natural application of the idea of exclusion statistics is offered by the various fractional quantum Hall effects. One may take the somewhat naive but certainly justifiable point of view that the essence of the $\nu = \frac{1}{m}$ fqHe is that under the appropriate conditions interacting electrons give rise to free quasi-particles with effective statistics parameter $g = m$. A familiar interpretation of these quasi-particles is that they can be viewed as composites of electrons plus an even number of flux quanta*. The familiar Laughlin wave functions describe the ground state configuration for these quasi-particles. The fundamental excitations (the Laughlin quasi-particles) are expected to carry the 'dual' (see section 3.3) statistics $g = \frac{1}{m}$.

The above scenario, which was suggested in Haldane's original paper, has been critically analyzed in the literature, where it has been confirmed in the appropriate low-temperature regime (see for example[16]). Our purpose in this paper is to set up and analyze a similar picture for *edge excitations* in the fqHe. Since such excitations can be described using the language of CFT, we first turn to a discussion of exclusion statistics in CFT spectra.

B. Exclusion statistics in CFT

Conformal Field Theories in two dimensions come with two commuting Virasoro algebras, and these infinite dimensional algebras can be used to organize the finite-size spectra of these theories. In such an approach, a CFT partition function is obtained by combining a number of characters of both Virasoro algebras (or extensions thereof). In applications such as String Theory, where the conformal symmetry has a geometric origin and the fundamental fields are bosonic coordinate fields, this 'Virasoro approach' to CFT is entirely natural. In contrast, the prototypical CFT in the condensed matter arena is a theory of free fermions, with a finite size spectrum that is simply a collection of many-fermion states constructed according to the rules set by the Pauli principle. When facing other CFT's that are relevant for condensed matter systems one may try to follow a similar road, which is to select a number of fundamental quasi-particle operators and to construct the full (chiral) spectrum as a collection of many-(quasi)-particle states. Explicit examples of this are the so-called spinon bases for a $\widehat{su(n)}_k$ Wess-Zumino-Witten models[20,21,18,22].

Let us now imagine that we have a concrete CFT, with explicit rules for the construction of a many-(quasi-)particle basis of the finite size spectrum. It is then natural to try to interpret that result in terms 'exclusion statistics' properties of the fundamental quasi-particles. In a recent paper[6], one of us has proposed a systematic procedure (based on recursion relations for truncated chiral spectra), which leads to 1-particle distribution functions for CFT quasi-particles. In many cases, it was established that the CFT thermodynamics are those of a free gas of quasi-particles governed by the new, generalized distribution functions. The examples discussed in[6] include spinons in the $\widehat{su(n)}_1$ WZW models, CFT parafermions and edge quasi-particles for the fractional qHe.

The example of the CFT for fqHe edge excitations is particularly interesting, since in those cases the generalized distributions derived from the CFT spectra are identical to those obtained from Haldane statistics (with specific values for g). In section 3 below we show in some detail how these results are established.

Clearly, the identification of Haldane statistics in fqHe edge theories is most useful since it provides a concrete link between rather abstract considerations on the systematics of quasi-particle bases on the one hand and concrete laboratory physics on the other. In particular, it opens up the possibility of analyzing transport phenomena such as edge-to-edge tunneling in the qHe (which has been well-studied both theoretically and experimentally) directly in terms of quasi-particles satisfying fractional exclusion statistics. We shall report the results of such an analysis in section 6 below.

III. QUASI-PARTICLES FOR THE $\nu = \frac{1}{M}$ FQHE EDGE

We consider the finite size spectrum for the CFT describing a single $\nu = \frac{1}{m}$ fqHe edge. In the CFT jargon, this theory is characterized as a $c = 1$ chiral free boson theory at radius $R^2 = m$. We shall consider the chiral Hilbert space corresponding to the following partition function

$$Z^{1/m}(q) = \sum_{Q=-\infty}^{\infty} \frac{q^{\frac{Q^2}{2m}}}{(q)_\infty} \ , \tag{III.1}$$

with $(q)_\infty = \prod_{l=1}^{\infty}(1-q^l)$ and $q = e^{-\beta \frac{2\pi}{L} \frac{1}{\rho_0}}$. [The 1-particle energies are of the form $\epsilon_l = l \frac{2\pi}{L} \frac{1}{\rho_0}$ with l integer and ρ_0 the density of states per unit length, $\rho_0 = (\hbar v_F)^{-1}$.] In this formula, the $U(1)$ affine Kac-Moody symmetry is clearly visible as all states at fixed $U(1)$ charge Q form an irreducible representation of this symmetry.

We should stress that the Hilbert space corresponding to (III.1) is not the physical Hilbert space for the edge theory of a quantum Hall sample with the topology of a disc. In the latter Hilbert space physical charge is quantized in units of e and, correspondingly, the $U(1)$ charge Q in (III.1) is restricted to multiples of m^4. In the geometry of a Corbino disc, *i.e.*, a cylinder, the operator that transfers charge $\frac{e}{m}$ from one edge to the other is physical. Accordingly, the physical Hilbert space is obtained by taking a tensor product of left and right copies of the Hilbert space (III.1) and restricting the total $U(1)$ charge $Q_L + Q_R$ to multiples of m^4. In the quasi-particle formalism that we present below the various restrictions on Q_L, Q_R are easily implemented.

Our goal here is to understand the collection of states in (III.1) in a different manner, and to view them as multi-particle states built from the creation operators for edge quasi-particles $\Psi_{\rm e}$ and $\Psi_{\rm qh}$. To simplify our notations, we shall write $G \equiv \Psi_{\rm e}$, $\phi \equiv {\Psi_{\rm qh}}^\dagger$. Due to the above-mentioned restrictions on the $U(1)$ charges Q and $Q_L + Q_R$, the chiral quasi-hole operator $\phi(z)$ by itself is not a physical operator in the edge theories for the disc or cylinder[‡] physical states are obtained by restricting the number of ϕ-quanta in the appropriate manner.

A. Quasi-hole states

We start by considering quasi-hole states that are built by applying only the modes ϕ_{-s} defined via $\phi(z) = \sum_s \phi_{-s} z^{s-\frac{1}{2m}}$. Clearly, the index s gives the dimensionless energy of the mode ϕ_{-s}. When acting on the charge-0 vacuum $|0\rangle$, we find the following multi-ϕ states (compare with[18] for the case $m = 2$, see also[23])

$$\phi_{-\frac{(2N-1)}{2m}-n_N} \cdots \phi_{-\frac{3}{2m}-n_2} \phi_{-\frac{1}{2m}-n_1} |\, 0\, \rangle \qquad \text{with} \quad n_N \geq n_{N-1} \geq \ldots \geq n_1 \geq 0 \; . \tag{III.2}$$

The choice of minimal modes is such that the lowest state of charge $Q\frac{e}{m}$ is at energy $\frac{Q^2}{2m}$, in agreement with the scaling dimension of the corresponding CFT primary field. Using so-called generalized commutation relations satisfied by the modes ϕ_{-s} one may show[18] that all multi-ϕ states different from (III.2) are either zero or linearly dependent on (III.2).

Before writing more general states we shall first focus on the exclusion statistics properties of the quanta ϕ_{-s}. We follow the procedure of[6] and start by introducing truncated partition sums for quasi-hole states (III.2). For $s = \frac{1}{2m}$, $\frac{3}{2m}$, etc, we define polynomials $P_s(x, q)$ to keep track of the number of many-body states that can be made using only the modes ϕ_{-k} with $k \leq s$, and that have a highest occupied mode with energy s' such that $s - s'$ is integer. $P_s(x, q)$ is defined as the trace of the quantity $x^N q^E$ over all these states, where N is the number of quasi-holes, E is the dimensionless total energy, and $x = e^{\beta\mu_{\rm qh}}$. For $m = 3$ this gives

$$P_{\frac{1}{6}} = xq^{\frac{1}{6}} \;, \quad P_{\frac{1}{2}} = x^2 q^{\frac{4}{6}} \;, \quad P_{\frac{5}{6}} = 1 + x^3 q^{\frac{9}{6}} \;, \quad \text{etc.} \tag{III.3}$$

In general, a occupied quasi-hole state of energy s corresponds to a factor xq^s in these generating polynomials.

The systematics of the edge quasi-hole states (III.2) directly lead to the following recursion relations between the polynomials $P_s(x, q)$,

$$P_s(x, q) = P_{s-1}(x, q) + x\, q^s\, P_{s-\frac{1}{m}}(x, q) \; . \tag{III.4}$$

For $m = 1$, which is the case corresponding to a $\nu = 1$ integer qHe edge, this relation directly implies $P_{l-\frac{1}{2}}(x, q) = \prod_{j=1}^{l} (1 + xq^{j-\frac{1}{2}})$. In that case the partition sum is simply a product and we recognize free fermions. For general m things are not that simple, but we can rewrite the recursion relation in matrix form

$$\begin{pmatrix} P_{l-\frac{2m-1}{2m}} \\ \vdots \\ P_{l-\frac{1}{2m}} \end{pmatrix} = M_l^{\rm qh}(x, q) \begin{pmatrix} P_{l-1-\frac{2m-1}{2m}} \\ \vdots \\ P_{l-1-\frac{1}{2m}} \end{pmatrix} , \tag{III.5}$$

with $l = 1, 2, \ldots$ and $M_l^{\rm qh}(x, q)$ the following $m \times m$ matrix

$$M_l^{\rm qh}(x, q) =$$

$$\begin{pmatrix} 1 & 0 & \ldots & 0 & xq^{l-\frac{2m-1}{2m}} \\ xq^{l-\frac{2m-3}{2m}} & 1 & \ldots & 0 & x^2 q^{2l-\frac{4m-4}{2m}} \\ \vdots & & \ddots & & \vdots \\ x^{m-1} q^{(m-1)l-\frac{m}{2}+\frac{2m-1}{2m}} & x^{m-2} q^{(m-2)l-\frac{m}{2}+\frac{4m-4}{2m}} & \ldots & xq^{l-\frac{1}{2m}} & 1 + x^m q^{ml-\frac{m}{2}} \end{pmatrix} . \tag{III.6}$$

The grand partition function for the quasi-hole states (III.2) is then given by

$$Z^{\rm qh}(x, q) = (\, 1 \quad 1 \quad \ldots \quad 1\,) \left(\prod_{l=1}^{\infty} M_l^{\rm qh}(x, q) \right) \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} . \tag{III.7}$$

We propose that the quasi-hole modes ϕ_{-s} with $s = l - \frac{2m-1}{2m}, \ldots, l - \frac{1}{2m}$ be viewed as a single (m-fold degenerate) level in the one-particle spectrum. [This convention is natural since a single quasi-particle over the ground state can only occupy one of these m levels.] The $m \times m$ matrix $M_l^{\rm qh}$ is then a level-to-level transfer matrix and replaces the free fermion ($m = 1$) factor $(1 + xq^{l-\frac{1}{2}})$. Clearly, the thermodynamics of the states (III.2) will be dominated by the largest eigenvalues $\lambda_l^+(x,q)$ of the matrices $M_l^{\rm qh}(x,q)$. These satisfy the characteristic equations

$$(\lambda_l^+ - 1)^m - x^m q^{ml-\frac{m}{2}} (\lambda_l^+)^{m-1} = 0 \ . \tag{III.8}$$

Instead of trying to solve these equations, we can derive from them a result for the 1-particle distribution functions

$$n^{\rm qh}(l) \equiv x\partial_x \ln(\lambda_l^+) = x \frac{\partial_x \lambda_l^+}{\lambda_l^+} \ . \tag{III.9}$$

We find

$$\begin{aligned} n^{\rm qh}(l) &= \frac{\lambda_l^+ - 1}{1 + \frac{1}{m}(\lambda_l^+ - 1)}, \\ (xq^l)^{-1} &= (\lambda_l^+ - 1)^{-1} (\lambda_l^+)^{1-\frac{1}{m}} \ . \end{aligned} \tag{III.10}$$

Comparing with (II.1) and identifying $g = \frac{1}{m}$ and $w(\epsilon) = (\lambda_l^+ - 1)^{-1}$, we see that the distribution function $n^{\rm qh}(l)$ becomes identical to $n_{g=\frac{1}{m}}(\epsilon = l)$. In other words, the exclusion statistics properties of the $\nu = \frac{1}{m}$ quasi-holes are those of 'ideal g-ons' in the sense of Haldane, with $g = \frac{1}{m}$! This identification is consistent with the result of bosonization applied to g-ons[24], and with the character computations of[25].

For the case $m = 2$, which is not in the category of fqHe edges, the equilibrium distribution is given by

$$n_{\frac{1}{2}}(\epsilon) = \frac{2}{\sqrt{1 + 4e^{-2\beta(\mu-\epsilon)}}} \ . \tag{III.11}$$

For $m = 3$ the explicit formulas (obtained using the Cardano formula for cubic equations) are quite unpleasant; figure 3.1 shows the distribution $n_{\frac{1}{3}}(\epsilon)$.



FIG. III.1. Distribution functions for fractional exclusion statistics with $g = 3$ (dashed line), $g = 1$ (dotted line), and $g = \frac{1}{3}$ (solid line) , all at the same temperature and at zero chemical potential.

B. Edge electron states

The same procedure can be applied to the edge electrons, which are created by modes G_{-t} with $G(z) = \sum_t G_{-t} z^{t-\frac{m}{2}}$. Multi-electron states take the form

$$G_{-(2M-1)\frac{m}{2}-m_M} \dots G_{-3\frac{m}{2}-m_2} G_{-\frac{m}{2}-m_1} | 0 \rangle \qquad \text{with} \quad m_M \geq m_{M-1} \geq \dots \geq m_1 \geq 0 \tag{III.12}$$

and we have truncated partition sums $Q_t(y,q)$ with t a half-odd-integer and $y = e^{\beta\mu_{\rm e}}$. They satisfy the recursion relations

$$Q_t(y,q) = Q_{t-1}(y,q) + y\, q^t\, Q_{t-m}(y,q), \tag{III.13}$$

with the following initial values

$$Q_{-\frac{m}{2}} = \dots = Q_{\frac{m}{2}-1} = 1 \ . \tag{III.14}$$

The 'transfer matrix' for the edge electrons $M_k^{\rm e}(y,q)$ is defined by

$$\begin{pmatrix} Q_K \\ \vdots \\ Q_{K+m-1} \end{pmatrix} = M_k^{\rm e}(y,q) \begin{pmatrix} Q_{K-m} \\ \vdots \\ Q_{K-1} \end{pmatrix} \tag{III.15}$$

with $K = km - m/2$, $k = 1, 2, \dots$ and we have

$$Z^{\rm e}(y,q) = (\, 1 \quad 0 \dots \quad 0 \quad 0\,) \left(\prod_{k=1}^{\infty} M_k^{\rm e}(y,q) \right) \begin{pmatrix} 1 \\ \vdots \\ 1 \\ 1 \end{pmatrix} . \tag{III.16}$$

In this case, a single action of the transfer matrix comprises a jump of m 1-particle levels, and the relevant distribution function will be

$$n^{\rm e}(k) \equiv y \partial_y \ln[(\mu_k^+)^{\frac{1}{m}}] = \frac{y}{m} \frac{\partial_y \mu_k^+}{\mu_k^+}, \tag{III.17}$$

The characteristic equation for the eigenvalue μ_k^+

$$\prod_{i=0}^{m-1} (\mu_k^+ - y q^{mk-i}) - (\mu_k^+)^{m-1} = 0 \ , \tag{III.18}$$

leads to

$$n^{\rm e}(k) = \frac{1}{m + (h_k - 1)} \ , \quad (y q^{mk})^{-1} = (h_k - 1)^m h_k^{1-m}. \tag{III.19}$$

with $h_k = \mu_k^+ / (y q^{mk})$. Identifying $w(\epsilon) = h_k - 1$, we again recognize the Wu equations (II.1) for Haldane exclusion statistics, this time with $g = m$, and we may identify $n^{\rm e}(k)$ with $n_{g=m}(\epsilon = mk)$.

For $m = 2$ this gives

$$n_2(\epsilon) = \tfrac{1}{2} \left(1 - \frac{1}{\sqrt{1 + 4\, e^{-\beta(\epsilon-\mu)}}} \right) \ . \tag{III.20}$$

See figure 3.1 for the distribution function $n_3(\epsilon)$ at $\mu_{\rm e} = 0$.

C. Duality

Having recognized distribution functions for fractional exclusion statistics with $g = \frac{1}{m}$ and $g = m$, respectively, we expect a particle-hole duality between the two cases (compare with[10,12]).

Before we come to that, we generalize the results of sections 3.1 and 3.2 by considering a chiral $c_{CFT} = 1$ CFT of compactification radius $R^2 = r/s$, with $r > s$ and r, s coprime. Choosing ϕ-quanta of charge $+\frac{s}{r} e$ and G-quanta of charge $-e$ as our fundamental excitations, we easily repeat the previous analysis and derive the following recursion relations

$$X_l(x) = X_{l-r}(x) + x\, X_{l-s}(x) \ , \qquad Y_l(y) = Y_{l-s}(y) + y\, Y_{l-r}(y) \ , \tag{III.21}$$

where we put $q = 1$ for convenience. [The connection with the quantities P_l and Q_l defined for $\frac{r}{s} = m$ is $X_l \leftrightarrow P_{\frac{2l-1}{2m}}$, $Y_l \leftrightarrow Q_{\frac{m}{2}+(l-1)}$.] Proceeding as before we obtain the distribution functions for Haldane statistics with $g = s/r$ (for the ϕ-quanta) and $\tilde{g} = r/s$ (for the G-quanta).

In the papers[10,12] it was recognized that the cases with g and $\tilde{g} = 1/g$ are dual in the sense that particles are dual to holes. To recover this duality in our present approach, we note that if $Y_l(y)$ is a solution of the second relation in (III.21), the expression

$$X_l(x) = Y_l(y = x^{-\tilde{g}})\, x^{\frac{l}{s}} \tag{III.22}$$

solves the first recursion relation. Assuming $r > s$, we can rewrite both recursion relations in a form involving a $r \times r$ recursion matrix. The largest eigenvalues $\lambda^+(x)$ and $\mu^+(y)$ are then related via

$$\lambda^+(x) = \mu^+(x^{-\tilde{g}})\, x^{\tilde{g}} \tag{III.23}$$

and the distribution functions

$$n_g(x) \equiv x\partial_x \ln \lambda^+(x) \ , \qquad n_{\tilde{g}}(y) \equiv g\, y\partial_y \ln \mu^+(y) \ , \tag{III.24}$$

satisfy

$$g\, n_g(x) = 1 - \tilde{g}\, n_{\tilde{g}}(y = x^{-\tilde{g}}) \ , \tag{III.25}$$

or, putting $\mu_{\tilde{g}} = -\tilde{g}\mu_g$ and restoring $q \neq 1$

$$g\, n_g(\epsilon) = 1 - \tilde{g}\, n_{\tilde{g}}(-\tilde{g}\epsilon) \ , \tag{III.26}$$

in agreement with the results of[10]. The interpretation of this result is that the $\tilde{g}$ quanta with positive energy act as holes in the ground state distribution of negative energy g-quanta. The relative factor $(-\tilde{g})$ between the energy arguments in (III.26) indicates that the act of taking out r g-quanta corresponds to adding s $\tilde{g}$-quanta. This duality further implies that, when setting up a quasi-particle description for fractional qHe edges, we can opt for (i) either quasi-holes òr edge electrons, with energies over the full range $-\infty < \epsilon < \infty$, or (ii) a combination of both types of quasi-particles, each having positive energies only. The CFT finite size spectrum naturally leads to (ii) (see section 3.4 below), while the analogy with Calogero-Sutherland quantum mechanics naturally leads to option (i) (see section 5). When considering transport equations in section 6, we shall be considering both alternatives.

D. The full spectrum

To complete our quasi-particle description for the $\nu = \frac{1}{m}$ edge, we need to specify how quasi-hole and electron operators can be combined to produce a complete basis for the chiral Hilbert space (III.1). We consider the following set of states

$$\begin{aligned} & G_{-(2M-1)\frac{m}{2}+Q-m_M} \ldots G_{-\frac{m}{2}+Q-m_1} \phi_{-(2N-1)\frac{1}{2m}-\frac{Q}{m}-n_N} \ldots \phi_{-\frac{1}{2m}-\frac{Q}{m}-n_1} |\, Q\, \rangle \\ & \text{with} \quad m_M \geq m_{M-1} \geq \ldots \geq m_1 \geq 0, \quad n_N \geq n_{N-1} \geq \ldots \geq n_1 \geq 0 \ , \\ & n_1 > 0 \quad \text{if} \quad Q < 0 \ , \end{aligned} \tag{III.27}$$

where $|Q\rangle$ denotes the lowest energy state of charge $Q\frac{e}{m}$ with Q taking the values $-(m-1)$, $-(m-2)$, …, -1, 0. Our claim is now that the collection (III.27) forms a basis of the chiral Hilbert space, so that

$$Z^{1/m}(q) = \sum_{Q=-(m-1)}^{0} q^{\frac{Q^2}{2m}}\, Z_Q^{\mathrm{qh}}(x=1,q)\, Z_Q^{\mathrm{e}}(y=1,q) \tag{III.28}$$

where we added a factor $q^{\frac{Q^2}{2m}}$ to take into account the energy of the initial states and we denoted by Z_Q^{qh} and Z_Q^{e} the generalizations of the partition functions (III.7) and (III.16) to the sector with vacuum charge Q. They are naturally written as

$$Z_Q^{\rm qh} = \sum_{N=0}^{\infty} \frac{q^{\frac{1}{2m}(N^2+2QN)+(1-\delta_{Q,0})N}}{(q)_N},$$

$$Z_Q^{\rm e} = \sum_{M=0}^{\infty} \frac{q^{\frac{m}{2}M^2-QM}}{(q)_M} \ , \tag{III.29}$$

with $(q)_L = \prod_{l=1}^{L}(1-q^l)$.

While the collection of states (III.27) looks rather complicated, it may be understood by considering the special case $m = 1$, which is a theory of two real free fermions of charge ±1. In this case there is only the $Q = 0$ vacuum and the allowed ϕ and G modes reduce to the familiar free fermion modes $\psi^{\pm}_{-\frac{1}{2}-n_j}$.

The right hand side of (III.28) has the form of a so-called 'fermionic sum formula'[17] and the equality of (III.1) and (III.28) is a new Rogers-Ramanujan identity. Similar identities relating 'fermionic sums' to characters in conformal field theories have been studied in the literature, see for example[17,18,19]. We would like to stress that the reasoning leading to these identities is very different between our approach and the work of[17]: in our approach the identities express exclusion statistics properties of CFT fields, while in the work of Kedem et al. the identities are based on Bethe Ansatz solutions of specific integrable lattice models. The first example where these two approaches have been explicitly connected is that of spinons in $SU(2)_1$ CFT and in the associated Haldane-Shastry spin chains[20,21,18].

The important conclusion from the above is that, up to a finite sum over vacuum charges, the chiral partition sum factorizes as a product of a quasi-hole piece and an edge electron piece. This means that the two types of quasi-particles are independent, or, in other words, that they do not have any mutual exclusion statistics. This then explains our asymmetric choice of quasi-particles. Had we chosen to work with fundamental quasi-particles of charges $\pm\frac{e}{m}$, we would have come across non-trivial mutual statistics. All of this is nicely illustrated with the case $m = 2$ where we can opt for the 'qHe basis' with independent quasi-holes and edge electrons, or for a 'spinon basis' built from charge $\pm\frac{e}{2}$ quanta, which are identical to the spinons of[20,21,18] and which have a non-trivial 2×2 statistics matrix. The two choices have the quasi-hole states (III.2) (called 'fully polarized spinon states' in[18]) in common, but differ in the way negative charges are brought in.

The observations made in this section may be generalized to composite edges such as those of the so-called Jain series with filling fraction $\nu = \frac{n}{np+1}$. We refer to appendix A for a brief discussion.

IV. EQUILIBRIUM QUANTITIES

A. Specific heat

The specific heat of a conformal field theory is well-known to be proportional to the central charge c_{CFT}

$$\frac{C(T)}{L} = \gamma \rho_0 k_B^2 T \ , \quad \gamma = \frac{\pi}{6} c_{CFT} \ , \tag{IV.1}$$

where $\rho_0 = (\hbar v_F)^{-1}$ is the density of states per unit length.

In[26] it was shown that the specific heat for g-on excitations, with energies in the full range $-\infty < \epsilon < \infty$, is in agreement with the central charge $c_{CFT} = 1$ of the corresponding CFT. The same result should of course come out in a picture where we select positive energy electrons and positive energy quasi-holes as our fundamental excitations. In this picture the total energy carried by the edge quasi-particles takes the form

$$E = \rho_0 \int_0^{\infty} d\epsilon \, \epsilon \, n_g(\epsilon) + \rho_0 \int_0^{\infty} d\epsilon \, \epsilon \, n_{\tilde{g}}(\epsilon) \ . \tag{IV.2}$$

and the corresponding result for the specific heat is

$$\frac{C(T)}{L} = (\gamma_{g,+} + \gamma_{\tilde{g},+}) \rho_0 k_B^2 T \ , \tag{IV.3}$$

where

$$\gamma_{g,+} = \partial_\beta \int_0^{\infty} d\epsilon \, \epsilon \, n_g(\epsilon) \ , \qquad \gamma_{\tilde{g},+} = \partial_\beta \int_0^{\infty} d\epsilon \, \epsilon \, n_{\tilde{g}}(\epsilon) \ . \tag{IV.4}$$

It takes an elementary application of the duality relation (III.26) to show that $\gamma_{\tilde{g},+} = \gamma_{g,-}$ and hence

$$(\gamma_{g,+} + \gamma_{\tilde{g},+}) = \gamma_g = \frac{\pi}{6} \ , \tag{IV.5}$$

confirming once again the value $c_{CFT} = 1$.

We would like to stress that the individual contributions $\gamma_{g,+}$ do depend on g and that only for $g = 1$ (Majorana fermions) $\gamma_{g,+}$ and $\gamma_{g,-}$ are equal. An exact result is[27]

$$\gamma_{g,+} = \frac{\pi}{6} \frac{L(\xi^g)}{L(1)} \ , \tag{IV.6}$$

with ξ a solution of the algebraic equation

$$\xi^g = 1 - \xi \tag{IV.7}$$

and $L(z)$ the Rogers dilogarithm. This gives

$$\gamma_{\frac{1}{2},+} = \frac{\pi}{6}\frac{3}{5}, \quad \gamma_{2,+} = \frac{\pi}{6}\frac{2}{5}, \quad \gamma_{\frac{1}{3},+} = \frac{\pi}{6} 0.655\ldots, \quad \gamma_{3,+} = \frac{\pi}{6} 0.344\ldots, \quad \text{etc.} \tag{IV.8}$$

B. Hall conductance

While the specific heat coefficient γ is not sensitive to g, the edge capacitance or, equivalently, the Hall conductance, obviously does depend on the filling fraction ν and thereby on g. In the quasi-particle formulation, this result comes out in a particularly elegant and simple manner.

Let us focus on a $\nu = \frac{1}{m}$ edge and take as our fundamental quasi-particles the edge electron of charge $q = -e$ and statistics $g = m$ and the edge quasi-hole of charge $q = \frac{e}{m}$ and statistics $\tilde{g} = \frac{1}{m}$, all quasi-particles having positive energies only.

Let us first consider zero temperature, where the Haldane distribution functions are step-functions with maximal value $n_g = \frac{1}{g}$. If we now put a voltage $V > 0$ the $q < 0$ quasi-particles will see their Fermi energy shift by the amount qV and all available states at energy up to $-qV$ will be filled. The total charge $\Delta Q(V, T = 0)$ that is carried by these excitations equals[§]

$$\Delta Q(V, T = 0) = \frac{1}{g} \cdot q \cdot \rho_0 \cdot (-qV) \tag{IV.9}$$

where the factor $\frac{1}{g}$ originates from the maximum of the distribution function and thus represents the statistics properties of the quasi-particles. Clearly, positive-q quasi-particles do not contribute to the response at $T = 0$, $V > 0$.

For the $\nu = \frac{1}{m}$ fqHe edges, the result for $V > 0$ is

$$\Delta Q(V > 0, T = 0) = \frac{1}{m} \cdot (-e) \cdot \rho_0 \cdot (eV) = -\frac{e^2}{m} \rho_0 V \tag{IV.10}$$

while for $V < 0$

$$\Delta Q(V < 0, T = 0) = m \cdot \frac{e}{m} \cdot \rho_0 \cdot (-\frac{e}{m} V) = -\frac{e^2}{m} \rho_0 V \ . \tag{IV.11}$$

Clearly, the edge capacitance

$$\frac{\Delta Q(V, T = 0)}{V} = -\rho_0 \frac{e^2}{m} \tag{IV.12}$$

is independent of the sign of V and we establish the correct value of the Hall conductance

$$G = \frac{1}{\rho_0 h} \frac{|\Delta Q|}{V} = \frac{1}{m} \frac{e^2}{h}. \tag{IV.13}$$

To show that the results (IV.12), (IV.13) hold for finite temperatures as well we write the general expression

$$\Delta Q(V,T) = -e\rho_0 \int_0^\infty d\epsilon \, n_m(\epsilon + eV) + \frac{e}{m}\rho_0 \int_0^\infty d\epsilon \, n_{\frac{1}{m}}(\epsilon - \frac{e}{m}V) \tag{IV.14}$$

and evaluate $\partial_\beta \Delta Q(V,T)$. Using once again the duality relations (III.26), we derive

$$\partial_\beta \Delta Q = \frac{e}{m}\,\rho_0 \partial_\beta \int_{-\infty}^\infty d\epsilon \, n_{\frac{1}{m}}(\epsilon) \propto \int_0^\infty dx \, \ln(x) \partial_x n_{\frac{1}{m}}(x) \tag{IV.15}$$

with $x = e^{-\beta\epsilon}$. Using (III.9) the last line turns into

$$\propto \lim_{x_0 \to \infty} \Big[\ln \lambda^+(x) - n_{\frac{1}{m}}(x) \ln(x) \Big]_0^{x_0} \tag{IV.16}$$

and by using the asymptotic behavior for $x \to \infty$

$$\lambda^+(x) \approx x^m \ , \qquad n_{\frac{1}{m}} \approx m \tag{IV.17}$$

we conclude that $\partial_\beta \Delta Q$ is indeed zero.

V. JACK POLYNOMIALS AND BEYOND

The quasi-particle basis that we specified in (III.27) has some arbitrariness to it. For example, we could have chosen to act first with the G_{-t} and then with ϕ_{-s}, which would have lead to a different set of states. Also, one quickly finds that the states (III.27) as they stand are not mutually orthogonal. For the purpose of establishing the thermodynamics of the fqHe edge theory, what matters is the counting of the number of states with given charge and energy, and this information can be extracted from (III.27). However, for the analysis of more detailed questions, in particular those concerning transport, the precise form of the multi-quasi-particle states is of crucial importance.

In this section, we shall present an ‘improved’ set of multi-particle states, which are mutually orthogonal and which are faithful to the statistics properties of the quasi-particles ϕ_{-s} and G_{-t}. The idea will be to specify an operator H_{CS} that acts on the CFT spectrum, and to modify the multi-particle states in such a way that they become eigenstates of H_{CS}. The operator H_{CS}, which was first given by Iso in[23], will be nothing else than a CFT version of the hamiltonian of so-called Calogero-Sutherland (CS) quantum mechanics with inverse square exchange. The analogy with CS quantum mechanics confirms the assignment of $g = \frac{1}{m}$ ($g = m$) exclusion statistics to ϕ_{-s} and G_{-t}, which are the CFT analogues of the particles and holes of the CS system. It also links the Jack polynomial eigenstates of the CS system to the quasi-particle basis of the fqHe edge theory.

We would like to stress that, in the context of the $\nu = \frac{1}{m}$ qHe edge, we do not assign physical significance to the operator H_{CS}. We merely use this operator as a device to select an optimal set of multi-particle states, where ‘optimal’ is meant in the sense of mutual orthogonality and of a relatively simple form of matrix elements of physical operators between the states.

The need for improving the form of the multi-particle states (III.27) can be phrased in yet another way. Let us, as an example, consider a multi-particle state containing two ϕ-quanta. If we were to work in position space, putting the two ϕ-fields at positions z_2 and z_1, the *exchange statistics* properties of the field $\phi(z)$ would result in simple phase factors associated to the interchange $z_2 \leftrightarrow z_1$ in a correlator. Working instead in energy space, with ϕ quanta ϕ_{-s_2} and ϕ_{-s_1}, we expect that the *exclusion statistics* properties of ϕ will imply simple behavior under the interchange $s_2 \leftrightarrow s_1$ in a correlator or form factor. In particular, one expects that interchanging $s_2 \leftrightarrow s_1$ in a form factor involving a state $|\ldots, s_2, s_1, \ldots\rangle$ will result in a phase factor $e^{i\frac{\pi}{m}}$. We shall show below (see, e.g., eq. V.10), that the form factors of the true ‘Jack polynomial’ multi-particle states $|\ldots, s_2, s_1, \ldots\rangle$ indeed satisfy this simple property, which is not valid for the naive multi-particle state $(\ldots \phi_{-s_2}\phi_{-s_1} \ldots)|0\rangle$. In mathematical terms, the issue is to define the correct co-product in a situation where, due to fractional statistics, the relevant symmetry is not a Lie algebra but rather a quantum group. In the context of the spinon basis for the $m = 2$ theory, this quantum group is a so-called yangian, and it has been established that the ‘Jack-polynomial’ co-product agrees with the co-product that is dictated by the quantum group symmetry[21,18].

A. The operator H_{CS}

To specify the operator H_{CS}, we employ the free boson $\varphi(z)$, which already featured in our formula (I.2). Following[23], we define

$$H_{CS} = \frac{m-1}{m} \sum_{l=0}^{\infty} (l+1)(i\sqrt{m}\partial\varphi)_{-l-1}(i\sqrt{m}\partial\varphi)_{l+1} + \frac{1}{3m}\left[(i\sqrt{m}\partial\varphi)^3\right]_0 \ , \tag{V.1}$$

where $\partial\varphi(z) = \sum_l (\partial\varphi)_l z^{-l-1}$ and where the second term on the r.h.s. denotes the zero-mode of the normal ordered product of three factors $(i\sqrt{m}\partial\varphi)(z)$. As a first result, one finds the following action of H_{CS} on states containing a single quasi-particle of charge $\frac{e}{m}$ or $-e$

$$H_{CS}\, \phi_{-\frac{1}{2m}-n}|0\rangle = h_\phi(n)\, \phi_{-\frac{1}{2m}-n}|0\rangle \ , \qquad \text{with} \qquad h_\phi(n) = \left[\frac{1}{3m} + mn(n+\frac{1}{m})\right]$$

and

$$H_{CS}\, G_{-\frac{m}{2}-n}|0\rangle = h_G(n) G_{-\frac{m}{2}-n}|0\rangle \ , \qquad \text{with} \qquad h_G(n) = \left[-\frac{m^2}{3} - n(n+m)\right] \ . \tag{V.2}$$

We would like to stress that the fact that both ϕ_s and G_t diagonalize H_{CS} is quite non-trivial. If one evaluates H_{CS} on any vertex operator $\phi^{(Q)}$ (of charge $Q\frac{e}{m}$), one typically runs into the field product $(T\phi^{(Q)})(z)$, where $T(z) = -\frac{1}{2}(\partial\varphi)^2(z)$ is the stress-energy of the scalar field φ. Only for $Q=1$ and $Q=-m$ do such terms cancel and do we find that the quasi-particle states are eigenstates of H_{CS}.

We can now continue and construct eigenstates of H_{CS} which contain several ϕ or G-quanta. What one then finds is that the simple product states such as (III.2) are not H_{CS} eigenstates, but that they rather act as head states that need to be supplemented by a tail of subleading terms. As an example, one finds two-ϕ eigenstates to be of the form[18]

$$|n_2, n_1\rangle = \phi_{-\frac{3}{2m}-n_2}\phi_{-\frac{1}{2m}-n_1}|0\rangle + \sum_{l=1}^{\infty} a_l \left[\phi_{-\frac{3}{2m}-n_2-l}\phi_{-\frac{1}{2m}-n_1+l}|0\rangle\right] \tag{V.3}$$

with coefficients a_l that can be computed. The connection of the coefficients a_l with the Jack polynomials that feature in the eigenfunctions in CS quantum mechanics has been made explicit in[18]. For the H_{CS} eigenstate headed by the multi-particle state (III.27) (with unit coefficient), we shall use the notation

$$|\{m_j\},\{n_i\}\rangle \tag{V.4}$$

so that

$$H_{CS}|\{m_j\},\{n_i\}\rangle = \left[\sum_{j=1}^{M} h_G((j-1)m + m_j) + \sum_{i=1}^{N} h_\phi(\frac{1}{m}(i-1) + n_i)\right] |\{m_j\},\{n_i\}\rangle \ . \tag{V.5}$$

Clearly, the states (V.4), with the m_j and n_i as specified in (III.27), form a complete and orthogonal basis for the chiral Hilbert space.

B. Norms and form factors

Of importance for later calculations are the norms of the states (V.4) and the matrix elements of physical operators between these states. For the explicit evaluation of such quantities we used the connection with Jack polynomials, relying on results that are available in the mathematical literature ([28], see also[29]).

As an example, we focus on multi quasi-hole states $|\{n_i\}\rangle$. To make contact with the Jack's, we view the ordered set $\{n_i\}$ as a Young tableau λ. The norm-squared of the state $|\{n_i\}\rangle$ then becomes

$$\langle\{n_i\}|\{n_i\}\rangle = j_{\lambda'} \tag{V.6}$$

where λ' is the Young tableau dual to λ and the j_λ are taken from[28]. Explicit examples are

$$\langle n_1|n_1\rangle = j_{(1^{n_1})} = \frac{(n_1 + \frac{1}{m} - 1)(n_1 + \frac{1}{m} - 2)\ldots\frac{1}{m}}{n_1(n_1-1)\ldots 1}$$

$$\langle n_2, n_1|n_2, n_1\rangle = j_{(2^{n_1},1^{n_2-n_1})}$$

$$= \frac{(n_2 + \frac{2}{m} - 1)(n_2 + \frac{2}{m} - 2) \dots (n_2 + \frac{2}{m} - n_1)}{(n_2 + \frac{1}{m})(n_2 + \frac{1}{m} - 1) \dots (n_2 + \frac{1}{m} - n_1 + 1)}$$

$$\times \frac{(n_2 - n_1 + \frac{1}{m} - 1) \dots \frac{1}{m}}{(n_2 - n_1) \dots 1} \, \frac{(n_1 + \frac{1}{m} - 1) \dots \frac{1}{m}}{n_1 \dots 1}$$

$$\text{etc.} \tag{V.7}$$

In the limit where all $n_i \gg 1$, one finds

$$\langle \{n_i\} | \{n_i\} \rangle \approx \prod_{i=1}^{N} \frac{{n_i}^{\frac{1}{m}-1}}{\Gamma(\frac{1}{m})} \ . \tag{V.8}$$

Of interest for the analysis of processes where electrons or holes tunnel into a fqHe edge is the form factor

$${}_N\langle \{n_m, \dots, n_2, n_1\} | \, G^\dagger_{-\frac{m}{2}-n} \, | \, 0 \, \rangle = f(n_m, \dots, n_1) \, \delta_{n, n_m + \dots + n_1} \ . \tag{V.9}$$

where the subscript N indicates that the state has been properly normalized. This form factor describes the amplitude by which an incoming hole (described by the operator $G^\dagger$ and of charge $+e$) creates a state that has m quasi-holes excited over the ground state. Explicit computation in the limit where all $n_i \gg 1$ yields (for simplicity we give the result for $m = 3$, see appendix B for the general case)

$$f(n_3, n_2, n_1) \approx \frac{\Gamma(\frac{1}{3})^{\frac{1}{2}}}{\Gamma(\frac{2}{3})} \left(\frac{(n_3 - n_2)(n_3 - n_1)(n_2 - n_1)}{n_3 \, n_2 \, n_1} \right)^{\frac{1}{3}} \ . \tag{V.10}$$

Remarkably, this result takes the form of a 'Jastrow factor' in the energy variables n_i. The order-($\frac{1}{3}$) zero's when two n_i come near reflect the $g = \frac{1}{3}$ exclusion statistics properties of the fundamental quasi-holes. Note that the expression (V.10) is invariant under global scalings of all energies n_i. The form (V.10) of the form factor can be viewed as a limit in (chiral) CFT of a result on correlation functions for the 'classical' model of quantum mechanics with inverse square exchange. This result was conjectured by Haldane[30] and later proven in[13,29].

VI. TRANSPORT PROPERTIES

Having checked that the thermodynamics of fqHe edges is correctly reproduced in the new quasi-particle language we are now ready to move on and consider transport properties. Following the set-up of a number of recent experiments, we shall consider a situation where electrons (or holes) from a fermi-liquid reservoir are allowed to tunnel into a $\nu = \frac{1}{m}$ fqHe edge. The DC *I*-*V* characteristic for this set-up, which were first computed in by Kane and Fisher[31] (see also[32]), show a cross-over from a linear (thermal) regime into a power-law behavior at high voltages and thus presents a clear fingerprint of the Luttinger liquid features of the fqHe edge. The experimental results from[8] are in agreement with these predictions. (See[2] for a further theoretical analysis of these data.)

The calculations by Kane and Fisher were based on bosonization and on the Keldysh formalism for non-equilibrium transport. Our goal here is to see if we can reproduce their results in an approach directly based on the edge quasi-particle formalism. Before going into this, we would like to stress that the 'Thermodynamic Bethe Ansatz (TBA) quasi-particles' behind the approach of[2] are quite different from what we have here, the most important distinction being that the TBA quasi-particles are a combination of degrees of freedom of both sides of the tunneling barrier; they do not exist for a $\nu = \frac{1}{m}$ edge in isolation.

If the $\nu = \frac{1}{m}$ fqHe edge were to behave as a fermi-liquid, we could calculate charge transport across a barrier using a simple (Boltzmann) kinetic equation of the form

$$I(V,T) \propto e \int_{-\infty}^{\infty} d\epsilon \, W \, \{ f_1(\epsilon - eV) F_2(\epsilon) - F_1(\epsilon - eV) f_2(\epsilon) \} \ , \tag{VI.1}$$

with $f(\epsilon)$ and $F(\epsilon)$ the Fermi-Dirac distributions for electrons and holes, respectively, and W the probability for an electron or hole of energy ϵ to cross the barrier and enter the edge. As is well known, this Boltzmann equation leads to an ohmic (linear in V) and temperature-independent current. Now that we have seen that the non-fermi liquid features of the $\frac{1}{m}$ edge can be captured via the statistics of the edge quasi-particles we can try to write a

'Boltzmann equation' for transport to and from fqHe edges by putting in appropriate generalizations $h(\epsilon)$ and $H(\epsilon)$ of the quantities $f_2(\epsilon)$ and $F_2(\epsilon)$, respectively. Before giving precise results (in section 6.1 below) we shall consider a 'naive' expression based on the intuition from the quasi-particle approach. In first approximation, the factor $h(\epsilon)$, which describes the probability for an electron to *leave* a $\nu = \frac{1}{m}$ edge, comprises two effects

1. a correlation effect, which can be traced to the non-trivial scaling dimension of the edge electron operator (see for example eq. (I.3)). At zero temperature, this is the so-called tunneling density of states

$$A^{+}(\epsilon) \propto \epsilon^{m-1} \; , \tag{VI.2}$$

2. a temperature dependence related to the exclusion statistics properties of the edge electrons. As we have seen, the natural factor associated to the *presence* of an edge electron is the distribution function

$$n_{g=m}(\epsilon) \; . \tag{VI.3}$$

Combining these factors, we come to the naive expressions

$$h^{(0)}(\epsilon) = \epsilon^{m-1}\, n_{g=m}(\epsilon) \; , \tag{VI.4}$$

and by similar reasoning we obtain

$$H^{(0)}(\epsilon) = \epsilon^{m-1}\, e^{\beta\epsilon}\, n_{g=m}(\epsilon) \; , \tag{VI.5}$$

where the thermal factor $e^{\beta\epsilon}\, n_m(\epsilon)$ has been dictated by the requirement of detailed balance.¶

One quickly finds that the Boltzmann equation with factors $h^{(0)}$ and $H^{(0)}$ is not exact at finite temperature. In section 6.2 we shall further comment on this equation and argue that it can be viewed as a first stage in a systematic approach. Before we come to that, we shall in the next section present a particularly simple derivation of the exact perturbative *I*-*V* characteristics for tunneling from a Fermi-liquid to a $\nu = \frac{1}{3}$ fqHe edge. This derivation uses the idea of a kinetic equation, together with the algebraic properties of the edge electrons.

A. Kinetic equation for inter edge transport

A careful derivation, based directly on the form of the tunneling hamiltonian‖

$$H_{int} \propto t \int d\epsilon \left[\Psi^{\dagger}_{\nu=1}(\epsilon) \Psi_{\nu=\frac{1}{3}}(\epsilon) + \text{h.c.} \right] \; , \tag{VI.6}$$

leads to the following kinetic equation (see e.g.[32])

$$I(V,T) \propto e\, t^2 \int_{-\infty}^{\infty} d\epsilon \, [f(\epsilon - eV) H(\epsilon) - F(\epsilon - eV) h(\epsilon)] \; , \tag{VI.7}$$

where h, H are one particle Green's functions

$$H(\epsilon) = \langle \Psi^{\dagger}_{\nu=\frac{1}{3}}(\epsilon) \Psi_{\nu=\frac{1}{3}}(\epsilon) \rangle_{V,T} \; , \qquad h(\epsilon) = \langle \Psi_{\nu=\frac{1}{3}}(\epsilon) \Psi^{\dagger}_{\nu=\frac{1}{3}}(\epsilon) \rangle_{V,T} \tag{VI.8}$$

for edge electrons in the $\nu = \frac{1}{3}$ fqHe edge, taken at $V = 0$. Note that the expression (VI.7) is perturbative as it gives the lowest non-trivial order in the parameter t.

The quantities $H(\epsilon)$ and $h(\epsilon)$ can be determined by using two simple observations. The first is that of *detailed balance*, which can be phrased as the requirement that at zero voltage there should be no current flowing. This fixes the ratio of $H(\epsilon)$ and $h(\epsilon)$ according to

$$H(\epsilon) = e^{\beta(\epsilon - eV)} h(\epsilon) \; . \tag{VI.9}$$

The second observation uses the *algebraic properties* of the edge electron operator, which include the anti-commutation relation

$$\left\{ \Psi^{\dagger}_{\nu=\frac{1}{3}}(\epsilon), \Psi_{\nu=\frac{1}{3}}(\epsilon') \right\} = \frac{2\pi}{L} \frac{1}{\rho_0} \epsilon^2 \delta(\epsilon - \epsilon') + 6 \frac{E}{\rho_0} + 3(\epsilon + \epsilon') \frac{\Delta Q}{e\rho_0} \; . \tag{VI.10}$$

In this formula, E is the operator for the total energy per unit length (proportional to the Virasoro zero mode L_0), and ΔQ is the operator for the total charge per unit length (proportional to the zero mode J_0 of the $U(1)$ Kac-Moody algebra). Clearly, this anti-commutator fixes the sum $H(\epsilon)+h(\epsilon)$. The expectation values of energy and charge follow directly from our analysis in section 4. We find

$$\langle E\rangle = \rho_0 \left(\frac{\pi^2}{6\beta^2}+\frac{(eV)^2}{6}\right) \ , \qquad \langle \Delta Q\rangle = -e\rho_0 \frac{(eV)}{3} \tag{VI.11}$$

and obtain the exact expressions

$$H(\epsilon) = \frac{(\epsilon-eV)^2+\frac{\pi^2}{\beta^2}}{e^{-\beta(\epsilon-eV)}+1} \ , \quad h(\epsilon) = \frac{(\epsilon-eV)^2+\frac{\pi^2}{\beta^2}}{1+e^{\beta(\epsilon-eV)}} \ . \tag{VI.12}$$

They lead to I-V characteristics

$$I(V,T) \propto e\, t^2\, \beta^{-3} \left(\frac{\beta eV}{2\pi}+\left(\frac{\beta eV}{2\pi}\right)^3\right) \ , \tag{VI.13}$$

in agreement with the result obtained in different approaches[31,2].

Clearly, the Green's functions (VI.8) can be evaluated in other ways, for example by using a conformal transformation in the x, t domain[32]. We would like to stress that our derivation is more direct and uses nothing more than the fundamental anti-commutation relation of the edge electrons. For $\nu = \frac{1}{3}$, these are particularly simple as they derive from the so-called $N = 2$ superconformal algebra, which has been well-studied in other contexts. For other filling fractions the fundamental anti-commutators look more complicated but are available in principle.

B. Interpretation in terms of exclusion statistics

If we compare the exact kinetic equation for $\nu = \frac{1}{3}$ with a naive generalized Boltzmann equation, we see that the mistake in the latter is in the approximation of the Green's function $h(\epsilon)$ by a the product $h^{(0)}(\epsilon)$ of a tunneling density of states times a Haldane distribution for fractional statistics.

The reason why this approximation turns out to be rather poor is that the operator $N(\epsilon) = \Psi^\dagger_{\nu=\frac{1}{3}}(\epsilon)\Psi_{\nu=\frac{1}{3}}(\epsilon)$ inside a fqHe edge is *not* to be viewed as a simple counting operator weighted by the appropriate power law of ϵ. This fact can be traced to the non-trivial operator terms (proportional to the energy and the charge operators) in the r.h.s. of (VI.10). To further illustrate this point we evaluated the expectation value of the operator $N(\epsilon)$ in a (normalized) one-electron state $|\epsilon'\rangle$

$$\langle \epsilon'|N(\epsilon)|\epsilon'\rangle \propto \epsilon^2\delta(\epsilon-\epsilon') + 6\frac{(\epsilon'-\epsilon)(\epsilon'^2+\epsilon^2)}{\epsilon'^2}\theta(\epsilon'-\epsilon) \ . \tag{VI.14}$$

This result shows an interaction effect in the action of $N(\epsilon)$ on a one-electron state: rather than just counting quanta of energy ϵ, the operator $N(\epsilon)$ is sensitive to the presence of quanta at energy $\epsilon' > \epsilon$ as well. In the Green's function $h(\epsilon)$ (for $\epsilon > 0$), the first term on the r.h.s. of (VI.14) corresponds to $h^{(0)}(\epsilon)$, while the second term leads to the following correction term

$$h^{(1)}(\epsilon) = 6\int_\epsilon^\infty d\epsilon' \, \frac{(\epsilon'-\epsilon)(\epsilon'^2+\epsilon^2)}{\epsilon'^2} \, n_3(\epsilon') \ . \tag{VI.15}$$

In figure 6.1 we have plotted the exact result for $h(\epsilon)$ against the approximations $h^{(0)}(\epsilon)$ and $\left[h^{(0)}+h^{(1)}\right](\epsilon)$. Clearly, the correction term $h^{(1)}(\epsilon)$ greatly improves the accuracy of the description.



FIG. VI.1. One-particle Green's function $h(\epsilon)$ for filling fraction $\nu = \frac{1}{3}$ and at zero voltage. The drawn curve is the exact result (VI.12); the dashed curve is the approximation $h^{(0)}(\epsilon)$and the dotted curve corresponds to $\left[h^{(0)} + h^{(1)}\right](\epsilon)$.

The situation here can be described as follows. As far as thermodynamics goes, the distribution functions $n_3(\epsilon)$ and $n_{\frac{1}{3}}(\epsilon)$ give exact results for quantities such as specific heat and conductance, and we may view the edge system as an ideal gas of fractional statistics quasi-particles. However, the operators $\Psi^\dagger_{\nu=\frac{1}{3}}(\epsilon)$, $\Psi_{\nu=\frac{1}{3}}(\epsilon)$ are not one-particle operators in the usual sense, as they do not simply add or extract a single quasi-particle from a many-particle state. In edge tunneling experiments, the edge system communicates with a Fermi liquid *via* the operators $\Psi^\dagger_{\nu=\frac{1}{3}}(\epsilon)$ and $\Psi_{\nu=\frac{1}{3}}(\epsilon)$ and we can not avoid interaction effects. We do believe, however, that a systematic expansion based on the quasi-particle picture is possible.

To avoid reference to a 'filled sea of negative energy edge electrons' we prefer to discuss transport in the picture where the fundamental quasi-particles are positive energy edge electrons and edge quasi-holes, respectively. If we we stick for a moment to the abovementioned 'naive' 0-th order approximation, we would arrive at the following lowest contribution to the tunneling current at voltage V and temperature T

$$\begin{aligned}
& I^{(0)}(V,T) \propto \\
& -e \int_0^\infty d\epsilon \, n_1(\epsilon - V)\, N_3(\epsilon)\, \epsilon^2 \\
& +e \int_0^\infty d\epsilon \, N_1(\epsilon - V)\, n_3(\epsilon)\, \epsilon^2 \\
& +e \int_0^\infty d\epsilon_3 \, d\epsilon_2 \, d\epsilon_1 \, n_1(\sum_i \epsilon_i + V)\, N_{\frac{1}{3}}(\epsilon_3) N_{\frac{1}{3}}(\epsilon_2) N_{\frac{1}{3}}(\epsilon_1) f^2(\epsilon_3, \epsilon_2, \epsilon_1) \\
& -e \int_0^\infty d\epsilon_3 \, d\epsilon_2 \, d\epsilon_1 \, N_1(\sum_i \epsilon_i + V) n_{\frac{1}{3}}(\epsilon_3) n_{\frac{1}{3}}(\epsilon_2) n_{\frac{1}{3}}(\epsilon_1) f^2(\epsilon_3, \epsilon_2, \epsilon_1)
\end{aligned} \qquad \text{(VI.16)}$$

where $f(\epsilon_3, \epsilon_2, \epsilon_1)$ is the form factor given in (V.10) and the integrations are over ordered sets of energies $\epsilon_3 \geq \epsilon_2 \geq \epsilon_1 \geq 0$. We have used the notation

$$N_g(\epsilon) = e^{\beta\epsilon} n_g(\epsilon) \; . \qquad \text{(VI.17)}$$

This result becomes exact in the limit $T \to 0$, where all distributions become step functions and the interaction effects disappear. Note that this formula has a clear asymmetry between electrons and holes: electrons that come into the edge settle as edge electrons, while incoming holes 'decay' into a total of three edge quasi-holes, with relative amplitudes given by the form factor $f(\epsilon_3, \epsilon_2, \epsilon_1)$. For $T = 0$, $V > 0$ the expression (VI.16) reduces to a single term

$$
\begin{aligned}
I^{(0)}(V>0,T=0) &\propto -e\int_{0\le\epsilon\le V} d\epsilon\,\epsilon^2 \\
&\propto V^3 \ , \qquad \text{(VI.18)}
\end{aligned}
$$

while $T=0$, $V<0$ it reduces to

$$
\begin{aligned}
I^{(0)}(V<0,T=0) &\propto e\int_{0\le\sum_i \epsilon_i\le V} d\epsilon_3\, d\epsilon_2\, d\epsilon_1\, f^2(\epsilon_3,\epsilon_2,\epsilon_1) \\
&\propto V^3 \ , \qquad \text{(VI.19)}
\end{aligned}
$$

In the latter case, the power law $I \propto V^3$ is a simple consequence of the fact that we perform three independent integrations $\int d\epsilon_3 d\epsilon_2 d\epsilon_1$ over quasi-hole energies, with a form factor $f(\epsilon_3,\epsilon_2,\epsilon_1)$ that is scale-invariant.

Clearly, the expression (VI.16) needs corrections. We believe that a systematic expansion, along the lines of the expansion $h(\epsilon)=h^{(0)}(\epsilon)+h^{(1)}(\epsilon)+\ldots$ that we have demonstrated above, is possible. We plan to demonstrate this in more detail in a future publication.

VII. CONCLUSIONS

The edge electrons that have been central in this paper are the edge analogues of the composite fermions (CF) used to describe bulk physics. We have made clear that, while the exchange statistics of these particles are fermionic, their exclusion statistics properties are not and are instead captured by non-trivial distribution functions $n_m(\epsilon)$ that take the place of the familiar Fermi-Dirac distribution. We have also investigated to what extent a quasi-particle picture, with edge electrons and edge quasi-holes as the fundamental quanta, can be used as a starting point for a quantitative analysis of transport. We have used algebraic properties of the $\nu=\frac{1}{3}$ edge electrons to derive exact results, and we have claimed that in general exclusion statistics properties may be used to set up a systematic expansion. In our view, these results hold some important lessons for other situations where fractional statistics quasi-particles have been proposed (spinons in $d=1$ quantum spin chains, anyons in $d=2$, etc.).

ACKNOWLEDGEMENTS

We thank A.W.W. Ludwig for many insightful comments and collaboration in the early stages of this project. KS thanks P. Bouwknegt, E. Fradkin, and Y.-S. Wu for discussions. Part of this work was done at the 1997 ITP Santa Barbara Workshop on 'Quantum Field Theory in Low Dimensions: from Condensed Matter to Particle Physics '. This research has further benefited from NATO Collaborative Research Grant SA.5-2-05(CRG.951303) and by support from the FOM foundation of the Netherlands.

APPENDIX A: COMPOSITE EDGES: JAIN SERIES

In this appendix we briefly describe a quasi-particle formulation of the composite edge theories corresponding to the filling fractions $\nu = \frac{n}{np+1}$ of the Jain series. These edge theories can be written as a collection of n free bosons, coupled via the topological K-matrix of the effective bulk Chern-Simons theory[4]. In[34] it was shown that the effective low-energy CFT for particles satisfying Haldane statistics with $n \times n$ statistical matrix G is a $c = n$ CFT with topological matrix $K = G^{-1}$. Inverting the argument we expect that the fundamental excitations of the CFT for qHe matrix K can be interpreted in terms of pseudo-particles satisfying fractional exclusion statistics with matrix $G = K^{-1}$.

An alternative and more natural approach to the Jain series edges would be to first perform a change of basis which separates a single charged mode from a set of $n-1$ neutral modes[35,36]. The latter are governed by an $\widehat{su(n)}_1$ affine Kac-Moody symmetry, and can be treated separately. An option is to view them as a set of n free parafermions in the sense of Gentile, see[6]. The CFT for the remaining charged mode is of the type that we described in this paper, with $g = \nu$. The entire edge theory is then described by a single (charged) g-on and a set of $\widehat{su(n)}_1$ degrees of freedom.

As an example of how the chiral Hilbert space works out, here is the example of $\nu = 2/5$, with K matrix

$$K = \begin{pmatrix} 3 & 2 \\ 2 & 3 \end{pmatrix} . \tag{A.1}$$

This theory has two independent $U(1)$ affine Kac-Moody symmetries, giving a factor $\left[\prod_{l=1}^{\infty}(1-q^l)\right]^{-2}$ in the partition function. The various charge sectors are labeled by pairs of integers (l_1, l_2), the energy being given by $E(l_1, l_2) = \frac{1}{10}(3l_1^2 - 4l_1 l_2 + 3l_2^2)$ (this is the bilinear form defined by the inverse of the K-matrix (A.1)). Thus

$$Z_{\nu=2/5}(q) = \sum_{(l_1,l_2)} \frac{q^{E(l_1,l_2)}}{\left[\prod_{l=1}^{\infty}(1-q^l)\right]^2}. \tag{A.2}$$

Under the rearrangement into $\widehat{su(2)}_1$ times $U(1)$, the combination $\frac{1}{2}(l_2 - l_1)$ plays the role of the $su(2)$ spin, while $l_1 + l_2$ is the charge under the new $U(1)$. The character identity will be[36]

$$Z_{\nu=2/5}(q) = \chi^{su(2)_1}_{j=0}(q) Z^{2/5}_{\text{even}}(q) + \chi^{su(2)_1}_{j=\frac{1}{2}}(q) Z^{2/5}_{\text{odd}}(q) , \tag{A.3}$$

where the subscript even (odd) on $Z^{s/r}$ means that we restrict to the states with total $U(1)$ charge Q even or odd. Simple expressions for the $su(2)_1$ characters are

$$\chi^{su(2)_1}_{j=0}(q) = \sum_{m+n \text{ even}} \frac{q^{\frac{1}{4}(m+n)^2}}{(q)_m (q)_n} , \qquad \chi^{su(2)_1}_{j=\frac{1}{2}}(q) = \sum_{m+n \text{ odd}} \frac{q^{\frac{1}{4}(m+n)^2}}{(q)_m (q)_n} . \tag{A.4}$$

For the general case with $\nu = \frac{n}{np+1}$, the charged sector is described by a free boson CFT at compactification radius $R^2 = \nu^{-1}$, which we write as $R^2 = \frac{r}{s}$. The chiral partition sum is

$$Z^{s/r}(q) = \sum_{Q=-\infty}^{\infty} \frac{q^{Q^2/(2rs)}}{\prod_{l=1}^{\infty}(1-q^l)} , \tag{A.5}$$

and restrictions, such as the even/odd in (A.3) are taken into account by restricting the charge quantum number Q.

Our fundamental charged edge quasi-particles will now be the primary fields of $U(1)$ charges $+s$ and $-r$; we shall write the creation and annihilation modes of these fields as ϕ_{-t} and G_{-t}, respectively. Note that for $s \neq 1$ the operators G_{-t} are not the physical edge electrons as the latter can only be written by including non-trivial factors from the neutral sector!

In close analogy with our analysis in section 3.4, we can now establish that the states

$$\begin{aligned} & G_{-(2M-1)\frac{r}{2s}+\frac{Q}{s}-m_M} \cdots G_{-\frac{r}{2s}+\frac{Q}{s}-m_1} \phi_{-(2N-1)\frac{s}{2r}-\frac{Q}{r}-n_N} \cdots \phi_{-\frac{s}{2r}-\frac{Q}{r}-n_1} | Q \rangle \\ & \text{with} \qquad m_M \geq m_{M-1} \geq \ldots \geq m_1 \geq 0, \quad n_N \geq n_{N-1} \geq \ldots \geq n_1 \\ & \text{and} \qquad n_1 \geq 0 \quad \text{if } Q \geq 0 \\ & \qquad\qquad n_1 > 0 \quad \text{if } Q < 0 , \end{aligned} \tag{A.6}$$

with $Q = -(r-s), \ldots, +(s-1)$, span the chiral Hilbert space (A.5) of the charged boson. The total energy of the lowest energy state in the charge sector Q having particle-numbers M and N for the quanta of type G and ϕ, respectively, equals

$$E(Q;M,N) = \frac{Q^2}{2rs} + \frac{r}{2s}M^2 - \frac{Q}{s}M + \frac{s}{2r}N^2 + \left[\frac{Q}{r} + \delta_{Q<0}\right] N \tag{A.7}$$

and this leads to the following expression for the chiral partition sum

$$Z^{s/r}(q) = \sum_{Q=-(r-s)}^{(s-1)} \sum_{M,N \geq 0} \frac{q^{E(Q;M,N)}}{(q)_M (q)_N} \; . \tag{A.8}$$

The equality of the expressions (A.5) and (A.8) is a new identity of the Rogers-Ramanujan type (see[17,18,19] for some similar identities).

In the case $p < 0$, the Jain series qHe edge exhibits counterflowing edge modes and it has been claimed that a new disorder-driven fixed point dominates the physics[37,35]. It will be most interesting to analyze this scenario in a quasi-particle formulation.

APPENDIX B: FORM FACTOR FOR GENERAL M

We briefly explain the exact evaluation of the form factor $f(m_m, \ldots, m_1)$ as defined in (V.9). Let us consider the special case $m = 2$ first. In that case the 'hole operator' $G(z)$ has conformal dimension 1 and may be identified with one of the currents of the affine Kac-Moody algebra $\widehat{su(2)}_1$. By exploiting the OPE

$$\phi(z)\phi(w) = (z-w)^{+\frac{1}{2}} \left[G(w) + \mathcal{O}(z-w) \right] \tag{B.1}$$

one obtains

$$G(w) = \oint_{C_w} \frac{dz}{2\pi i} (z-w)^{-\frac{3}{2}} \phi(z)\phi(w) \; . \tag{B.2}$$

We also have[21]

$$\phi(z)\phi(w)|\,0\,\rangle = (z-w)^{\frac{1}{2}} \sum_{n_2,n_1} P^{(-\frac{1}{2})}_{n_2,n_1}(z,w)|n_2,n_1\rangle \; , \tag{B.3}$$

where $P^{(-\frac{1}{2})}_{\{n_i\}}(z,w)$ are the appropriate Jack polynomials. Combining the above, we obtain

$$G(w)|\,0\,\rangle = \sum_{n_2,n_1} P^{(-\frac{1}{2})}_{n_2,n_1}(w,w)|n_2,n_1\rangle \tag{B.4}$$

and it follows that

$${}_N\langle n_2,n_1|\,G^\dagger_{-1-n}\,|\,0\,\rangle = \delta_{n,n_2+n_1} \; \left[j_{(2^{n_1},1^{n_2-n_1})} \right]^{\frac{1}{2}} \; P^{(-\frac{1}{2})}_{n_2,n_1}(1,1) \; , \tag{B.5}$$

with $j_{\lambda'}$ as in (V.6). For general m one obtains a similar result in terms of Jack polynomials with label $(-\frac{1}{m})$. Using the explicit result[28,29]

$$P^{(-\frac{1}{m})}_{\{n_i\}}(1,1) = \prod_{i=0}^{m-1} \frac{\Gamma(\frac{1}{m})}{\Gamma(1-\frac{i}{m})} \prod_{i<j} \frac{\Gamma(n_j - n_i + \frac{j-i+1}{m})}{\Gamma(n_j - n_i + \frac{j-i}{m})} \; , \tag{B.6}$$

together with the result (V.8) for the $j_{\lambda'}$, we derive the following asymptotic form for $n_i >> 1$

$$f(n_m, \ldots, n_1) = \frac{[\Gamma(\frac{1}{m})]^{m/2}}{\prod_{i=0}^{m-1} \Gamma(1-\frac{i}{m})} \frac{\prod_{i<j} (n_j - n_i)^{\frac{1}{m}}}{\prod_{i=1}^{m} n_i^{\frac{(m-1)}{2m}}} \; . \tag{B.7}$$

The simple Jastrow form of this form factor is a clear indication that in the limit $n_i \gg 1$ a much simpler derivation, along the lines of 'bosonization in momentum space' should be possible.

---

[1] P. Fendley, A.W.W. Ludwig and H. Saleur, Phys. Rev. Lett. **74**, 3005 (1995) ; Phys. Rev. **B52**, 8934 (1995) .

[2] C. de Chamon and E. Fradkin, Phys. Rev. **B56**, 2012 (1997).

[3] N.P. Sandler, C. de Chamon and E. Fradkin, Phys. Rev. **B57**, 12324 (1998)

[4] X.G. Wen, Phys. Rev. **B41**, 12838 (1990); Int. Jour. Mod. Phys. **B6**, 1711 (1992).

[5] M. Stone and M.P.A. Fisher,Int. J. Mod. Phys. **B8**, 2539 (1994)

[6] K. Schoutens, Phys. Rev. Lett. **79**, 2608 (1997).

[7] F.D.M. Haldane, Phys. Rev. Lett. **67**, 937 (1991).

[8] A.M. Chang, L.N. Pfeiffer and K.W. West, Phys. Rev. Lett. **77**, 2538 (1996).

[9] Y.-S. Wu, Phys. Rev. Lett. **73**, 922 (1994).

[10] C. Nayak and F. Wilczek, Phys. Rev. Lett. **73**, 2740 (1994).

[11] S.B. Isakov, Mod. Phys. Lett. **B8**, 319 (1994).

[12] A.K. Rajagopal, Phys. Rev. Lett. **74**, 1048 (1994).

[13] Z. Ha, Phys. Rev. Lett. **73**, 1574 (1994).

[14] M.V.N. Murthy and R. Shankar, Phys. Rev. Lett. **73**, 3331 (1994).

[15] A. Dasnières de Veigy and S. Ouvry, Phys. Rev. Lett. **72**, 600 (1994).

[16] K. Tevosyan ans A.H. MacDonald, *Virial Expansions, Exclusion Statistics and the Fractional Quantum Hall Effect*, preprint cond-mat/9701178; Y.S. Wu, Y. Yu, Y. Hatsugai and M. Kohmoto, Phys. Rev. **B57**, 9907 (1998)

[17] R. Kedem, T.R. Klassen, B.M. McCoy and E. Melzer, Phys. Lett. **B304**, 263 (1993); Phys. Lett. **B307**, 68 (1993).

[18] P. Bouwknegt, A.W.W. Ludwig and K. Schoutens, Phys. Lett. **B338**, 448 (1994); in Proceedings of the 1994 Trieste Summer School on High Energy Physics and Cosmology, E. Gava et al. eds (World Scientific 1995).

[19] P. Bouwknegt, A.W.W. Ludwig and K. Schoutens, Phys. Lett. **B359**, 304 (1995).

[20] F.D.M. Haldane, Phys. Rev. Lett. **66**, 1529 (1991); F.D.M. Haldane, Z.N.C. Ha, J.C. Talstra, D. Bernard and V. Pasquier,*ibid.* **69**, 2021 (1992).

[21] D. Bernard, V. Pasquier and D. Serban, Nucl. Phys. **B428**, 612 (1994).

[22] P. Bouwknegt and K. Schoutens, Nucl. Phys. **B482**, 345 (1996).

[23] S. Iso, Nucl. Phys. **B443**, 581 (1995).

[24] Y.-S. Wu and Y. Yu, Phys. Rev. Lett. **75**, 890 (1995).

[25] K. Hikami, Phys. Lett. **A205**, 364 (1995).

[26] S.B. Isakov, D.P. Arovas, J. Myrheim and A.P. Polychronakos, Phys. Lett. **A212**, 299 (1996).

[27] W. Nahm, A. Recknagel and M. Terhoeven, Mod. Phys. Lett. **A8**, 1835 (1993).

[28] R.P. Stanley, Adv. Math. **77**, 76 (1989).

[29] F. Lesage, V. Pasquier and D. Serban, Nucl. Phys. **B435**, 585 (1995).

[30] F.D.M. Haldane, in Proceedings of the 16th Taniguchi Symposium, Kashikojima, Japan, October 1993, A. Okiji and N. Kawakami, eds. (Springer Verlag 1994).

[31] C.L. Kane and M.P.A. Fisher, Phys. Rev. Lett. **68** (1992) 1220; Phys. Rev. **B46**, 15233 (1992).

[32] X.G. Wen, Phys. Rev. **B44**, 5708 (1991).

[33] R.K. Bhaduri, R.S. Bhalerao and M.V.N. Murthy, J. of Stat. Phys. **82**, 1659 (1996).

[34] T. Fukui and N. Kawakami, Phys. Rev. **B51**, 5239 (1995).

[35] C.L. Kane and M.P.A. Fisher, Phys. Rev. **B51**, 13449 (1995).

[36] A. De Martino and R. Musto, Int. Jour. Mod. Phys. **B9**, 2839 (1995).

[37] C.L. Kane, M.P.A. Fisher and J. Polchinski, Phys. Rev. Lett. **72**, 4129 (1994).

[*] The familiar terminology 'composite fermions' is somewhat unfortunate in this context, as we argue that the *exclusion* statistics properties of these composite quasi-particles are *not* fermionic. Clearly, their *exchange* statistics, which are determined 'modulo 2π', are fermionic.

[†] The notation G is inspired by the fact that the fundamental anti-commutation relations for the modes G_{-t} at $\nu = 1/3$ are those of the so-called $N = 2$ superconformal algebra. See section 6 for more on this.

[‡] In the proper mathematical terminology we call $\phi(z)$ a Chiral Vertex Operator (CVO).

[§] We use the symbol ΔQ for the total charge, while we keep Q for the reduced charge (in units of $\frac{e}{m}$).

[¶] This same thermal factor was proposed in the paper[33], which proposes a generalization of the Boltzmann equation to the case of fractional statistics. In our view, the proposal of[33] is incomplete, as it ignores the correlation effects which are unavoidable for quasi-particles obeying fractional exclusion statistics.

[‖] We write Ψ for the edge electron operator, (denoted by Ψ_e in section 2 and by G in section 3), and we indicate the filling fraction by an explicit subscript.